

Diffraction at small M^2/Q^2 in the QCD dipole picture

A. Bialas^{a,b}, H. Navelet^a and R. Peschanski^a

^aService de Physique Theorique CEA-Saclay
F-91191 Gif-sur-Yvette Cedex, France

^bM.Smoluchowski Institute of Physics, Jagellonian University
Reymonta 4, 30-059 Cracow, Poland

February 7, 2008

Abstract

Using the QCD dipole picture of BFKL dynamics, the quasi elastic component of the diffractive γ^* -dipole cross-section, dominating at small M^2/Q^2 , is calculated.

1. Diffraction dissociation of virtual photons, a process being on the border line between the perturbative and non-perturbative regions, represents a challenging problem for Quantum Chromodynamics [1]. It is therefore of no surprise that it became a subject of numerous studies by various methods [2]-[12].

The present investigation continues our effort [6]-[8] to reach a unified description of both total and diffractive cross-sections of virtual photons in the framework of the QCD dipole picture of high-energy interactions [9, 10]. More precisely, it is an attempt to obtain -within the approximations inherent to the QCD dipole model - an exact formula for the quasi-elastic component [6] of the (virtual) photon diffractive dissociation at arbitrary momentum transfer. It is a generalization of a recent calculation [8] of the forward photon diffractive processes. The need for an *exact* formula is emphasized by (i) recent success of the approximate dipole model description

[11, 12] of the HERA data [13] and (ii) improved accuracy of recent measurements [14] which provide already first results on momentum transfer dependence of diffractive structure functions.

We restrict ourselves to the so called quasi elastic component of the diffractive cross-section, dominating at small M^2/Q^2 where M is the mass of the produced diffractive system and Q is the virtuality of the photon. The calculation of the high mass component at non-zero momentum transfer was already presented in [7].

In the QCD dipole picture the cross-section for the quasi-elastic scattering of a virtual photon on a single dipole target can be written as [6]

$$\frac{d\sigma}{dM^2 d^2 p_T} = \frac{2N_c}{4\pi^2} \int d^2 k |< \vec{k}, M^2 | T_{p_T}^{qel} | Q >|^2 \quad (1)$$

where p_T is the transverse momentum after the collision, $\vec{k} = (|k|, \phi_k)$ is the relative tranverse momentum in the $(q\bar{q})$ pair, and

$$< \vec{k}, M^2 | T_{p_T}^{qel} | Q > = \int_0^1 \int d^2 r < \vec{k}, M^2 | \vec{r}, z > < \vec{r}, z | T_{p_T} | \vec{r}, z > \Psi(\vec{r}, z; Q). \quad (2)$$

Here $\vec{r} = (|r|, \phi_r)$ is the relative transverse distance in the $(q\bar{q})$ pair, z is the light-cone momentum fraction of one of the quarks and $< \vec{r}, z | T_{p_T} | \vec{r}, z >$ is the elastic dipole amplitude for scattering of the dipole of tranverse size r on the target dipole of size r_0 . $\Psi(\vec{r}, z; Q)$ are the light-cone photon wave functions [15].

Following the argument of [6] (c.f. also [8]), Eq.(1) can be rewritten in the form

$$\frac{d\sigma}{dM^2 d^2 p_T} = \frac{2N_c}{4\pi^2} \int_0^1 dz z(1-z) \frac{1}{2} d\phi_k \left| G(\hat{M}, z, p_T; x_{\mathcal{P}}) \right|^2. \quad (3)$$

where

$$G(k, z, p_T; x_{\mathcal{P}}) = \frac{1}{2\pi} \int d^2 r e^{i\hat{M}\vec{r}} < \vec{r}, z | T_{p_T} | \vec{r}, z > \Psi(r, z; \hat{Q}) \quad (4)$$

and $\hat{M}$ is the vector of length

$$\hat{M} = M\sqrt{z(1-z)} \quad (5)$$

parallel to $\vec{k}$. Eq.(4) is the starting point of our calculation.

2. The difficult part of the task is to perform the integration over d^2r in (4). To this end we need an adequate formula for the dipole-dipole amplitude $\langle \vec{r}, z | T_{p_T} | \vec{r}, z \rangle$. For $p_T = 0$ an explicit formula is available [9] and this allowed the calculation of [8]. For $p_T \neq 0$, however, the calculation is much more involved. In the present paper we perform it using the methods developed recently in [16].

The starting point is the general expression [17]

$$\langle \vec{r}, z | T_{p_T} | \vec{r}, z \rangle = 4\pi\alpha^2 \sum_{n=-\infty}^{n=\infty} \int \frac{d\nu}{\pi} e^{\omega(n,\nu)Y} d_{n,\nu} |r| E_{p_T}^{n,\nu}(r) |r_0| \bar{E}_{p_T}^{n,\nu}(r_0), \quad (6)$$

where

$$d_{n,\nu} = \left(16 \left[\nu^2 + \left(\frac{n-1}{2} \right)^2 \right] \left[\nu^2 + \left(\frac{n+1}{2} \right)^2 \right] \right)^{-1}, \quad (7)$$

$$\omega(n,\nu) = \frac{2\alpha N_c}{\pi} \left(\psi(1) - \text{Re}[\psi(\frac{1}{2}(|n|+1) + i\nu)] \right) \quad (8)$$

and $E_{p_T}^{n,\nu}(r)$ are eigenfunctions of the conformal operator in the mixed representation defined in [17]. The explicit expression for $E_{p_T}^{n,\nu}(r)$ is fairly complicated [16, 17] but, fortunately, we shall not need it here. We only quote, for future reference, the formula for $E_0^{n,\nu}(r)$:

$$E_0^{n,\nu}(r) = |r|^{-2i\nu} e^{in\phi_r}. \quad (9)$$

The sum over n in (6) was introduced for technical reasons. In the high-energy (large Y) limit, the term with $n = 0$ dominates.

To proceed, it is useful at this point to introduce the explicit formulae for $\Psi(\vec{r}, z; Q)$. We write them in the form

$$\Psi(\vec{r}, z; Q) = C \hat{Q} \Phi_{T,L}(z) \chi_{T,L}(r, \hat{Q}) \quad (10)$$

with $\hat{Q} = Q\sqrt{z(1-z)}$, $C = \sqrt{\alpha_{em}e_{(f)}}/2\pi$ ($e_{(f)}$ is the charge of a quark) and

$$\Phi_T = z; \quad \Phi_L = 2\sqrt{z(1-z)}; \quad \chi_T = e^{i\phi_r} K_1(\hat{Q}r); \quad \chi_L = K_0(\hat{Q}r). \quad (11)$$

The subscripts (T, L) denote transverse (right-handed) and longitudinal polarizations of the incident photon. Using (10) one obtains

$$G_{T,L}(k, z, p_T; x_{\mathcal{P}}) = D |r_0| \hat{Q} \Phi_{T,L}(z) \sum_n \int \frac{d\nu}{\pi} d_{n,\nu} e^{\omega(n,\nu)Y} E_{p_T}^{n,\nu}(r_0) g_{T,L}^{n,\nu}, \quad (12)$$

where $D \equiv 4\pi\alpha^2 C = 2\alpha^2 \sqrt{\alpha_{em}} e_{(f)}$,

$$g_{T,L}^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{2\pi} \int d^2r \psi_{T,L}(r, \hat{Q}, \hat{M}) E_{p_T}^{n,\nu}(r) \quad (13)$$

and

$$\psi_{T,L}(r, \hat{Q}, \hat{M}) = e^{i\hat{M}r} \chi_{T,L}(r, \hat{Q}) |r|. \quad (14)$$

We can now expand $\psi_{T,L}(r, \hat{Q}, \hat{M})$ in terms of conformal eigenfunctions at $p_T = 0$:

$$\psi_{T,L}(r, \hat{Q}, \hat{M}) = \sum_n \int d\nu \psi_{T,L}^{n,\nu}(\hat{Q}, \hat{M}) \frac{E_0^{n,\nu}(r)}{|r|^2} \quad (15)$$

The inverse tranform is

$$\psi_{T,L}^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{2\pi^2} \int d^2r \psi_{T,L}(r, \hat{Q}, \hat{M}) E_0^{n,\nu}(r) \quad (16)$$

and thus we obtain

$$g_{T,L}^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{2\pi} \sum_{n'} \int d\nu' \psi_{T,L}^{n',\nu'}(\hat{Q}, \hat{M}) \mathcal{I}_{p_T}^{n,n',\nu,\nu'}. \quad (17)$$

We see that the integration over r is reduced to

$$\mathcal{I}_{p_T}^{n,n',\nu,\nu'} \equiv \int d^2r \frac{E_{p_T}^{n,\nu}(r) E_0^{n',\nu'}(r)}{|r|^2} \quad (18)$$

which was explicitly calculated in [16] with the result

$$\begin{aligned} \mathcal{I}_{p_T}^{n,n',\nu,\nu'} &= \frac{\pi}{2} (-1)^{\frac{n-n'}{2}} \left[\frac{p_T}{8} \right]^{\tilde{\mu}-\tilde{\mu}'} \left[\frac{\tilde{p}_T}{8} \right]^{\mu-\mu'} \\ &\times \frac{\Gamma(1-\mu)}{\Gamma(\tilde{\mu})} \frac{\Gamma\left(\frac{\mu+\mu'}{2}\right) \Gamma\left(\frac{-\mu+\mu'}{2}\right)}{\Gamma\left(1-\frac{\tilde{\mu}+\tilde{\mu}'}{2}\right) \Gamma\left(1-\frac{\tilde{\mu}'-\tilde{\mu}}{2}\right)}, \end{aligned} \quad (19)$$

for $n - n'$ even and 0 for $n - n'$ odd. Here $\mu = i\nu - \frac{n}{2}$; $\tilde{\mu} = i\nu + \frac{n}{2}$.

We can now calculate $\psi_{T,L}^{n,\nu}(\hat{Q}, \hat{M})$. From (16) and (9) we have

$$\psi_{T,L}^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{2\pi^2} \int d^2r \psi_{T,L}(r, \hat{Q}, \hat{M}) |r|^{-2i\nu} e^{in\phi_r}, \quad (20)$$

where ϕ_r is the r azimuthal angle with respect to p_T .

Using

$$\int d\Psi e^{i\hat{M}\rho \cos \Psi \pm im\Psi} \equiv 2\pi e^{im\frac{\pi}{2}} J_m(\hat{M}\rho), \quad (21)$$

we find for right-handed photons (for left-handed photons, $n + 1$ should be replaced by $n - 1$):

$$\psi_T^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{\pi} \int r^2 dr r^{-2i\nu} e^{i(n+1)\frac{\pi}{2} - i\phi_k} J_{n+1}(\hat{M}r) K_1(\hat{Q}r). \quad (22)$$

For the longitudinal photons we obtain

$$\psi_L^{n,\nu}(\hat{Q}, \hat{M}) = \frac{1}{\pi} \int r^2 dr r^{-2i\nu} e^{in\frac{\pi}{2}} J_n(\hat{M}r) K_0(\hat{Q}r) \quad (23)$$

Using now [18]

$$\int_0^\infty d\rho \rho^{-\lambda} K_h(\hat{Q}\rho) J_l(\hat{M}\rho) = \frac{1}{4} \left(\frac{M}{Q}\right)^l \left(\frac{\hat{Q}}{2}\right)^{\lambda-1} \times \frac{\Gamma\left(\frac{1-\lambda+l+h}{2}\right) \Gamma\left(\frac{1-\lambda+l-h}{2}\right)}{\Gamma(l+1)} {}_2F_1\left(\frac{1-\lambda+l+h}{2}, \frac{1-\lambda+l-h}{2}, l+1; -\frac{M^2}{Q^2}\right) \quad (24)$$

and

$$\begin{aligned} & {}_2F_1\left(\frac{1-\lambda+l+h}{2}, \frac{1-\lambda+l-h}{2}, l+1; -\frac{M^2}{Q^2}\right) = \\ & = (\beta)^{\frac{1-\lambda+l-h}{2}} {}_2F_1\left(\frac{1-\lambda+l-h}{2}, \frac{1+\lambda+l-h}{2}, l+1; 1-\beta\right) \end{aligned} \quad (25)$$

where h, l are positive integers, we find

$$\begin{aligned} \psi_{T,right}^{n,\nu}(\hat{Q}, \hat{M}) &= \frac{1}{4\pi} e^{-i\pi\frac{n+1}{2} - i\phi_k} \left(\frac{M}{Q}\right)^{|n+1|} \left(\frac{\hat{Q}}{2}\right)^{2i\nu-3} \times \\ & \frac{\Gamma(-i\nu+2+\frac{|n+1|}{2}) \Gamma(-i\nu+1+\frac{|n+1|}{2})}{\Gamma(|n+1|+1)} (\beta)^{-i\nu+1+\frac{|n+1|}{2}} \times \\ & {}_2F_1\left(-i\nu+1+\frac{|n+1|}{2}, \frac{|n+1|}{2}+i\nu-1, |n+1|+1; 1-\beta\right), \end{aligned} \quad (26)$$

$$\begin{aligned}
\psi_L^{n,\nu}(\hat{Q}, \hat{M}) &= \frac{1}{4\pi} e^{-i\pi\frac{n}{2} - i\phi_k} \left(\frac{M}{Q}\right)^{|n|} \left(\frac{\hat{Q}}{2}\right)^{2i\nu-3} \times \\
&\quad \frac{\Gamma^2\left(-i\nu + \frac{3}{2} + \frac{|n|}{2}\right)}{\Gamma(|n|+1)} (\beta)^{-i\nu + \frac{3}{2} + \frac{|n|}{2}} \times \\
&\quad {}_2F_1\left(-i\nu + \frac{3}{2} + \frac{|n|}{2}, \frac{|n|}{2} + i\nu - \frac{1}{2}, |n|+1; 1-\beta\right). \tag{27}
\end{aligned}$$

Next we denote

$$\begin{aligned}
r_T(\nu_1, \nu_2) &= 2B(-i\nu_2 + i\nu_1 + 2, -i\nu_2 + i\nu_1) \\
r_L(\nu_1, \nu_2) &= 4B(-i\nu_2 + i\nu_1 + 1, -i\nu_2 + i\nu_1 + 1), \tag{28}
\end{aligned}$$

which are factors coming from integrals over the variable z . All in all, we obtain

$$\begin{aligned}
\frac{d\sigma}{dM^2 d^2 p_T} &= \frac{N_c}{2\pi} \int_0^1 dz \, z(1-z) \left| G(\hat{M}, z; x_{\mathcal{P}}) \right|^2 \\
&= \frac{N_c}{8\pi} Q^2 D^2 \sum_{n_1} \int \frac{d\nu_1}{\pi} e^{\omega(n_1, \nu_1)} Y_{d_{n_1, \nu_1}} |r_0| \bar{E}_{p_T}^{n_1, \nu_1}(r_0) \\
&\quad \times \sum_{n_2} \int \frac{d\nu_2}{\pi} e^{\omega(n_2, \nu_2)} Y_{\bar{d}_{n_2, \nu_2}} |r_0| E_{p_T}^{n_2, \nu_2}(r_0) \\
&\quad \times \sum_{n'_1} \int \frac{d\nu'_1}{\pi} \psi_{T,L}^{n'_1, \nu'_1}(Q, M) \mathcal{I}_{P_T}^{n_1, n'_1, \nu_1, \nu'_1} \\
&\quad \times \sum_{n'_2} \int \frac{d\nu'_2}{\pi} \bar{\psi}_{T,L}^{n'_2, \nu'_2}(Q, M) \bar{\mathcal{I}}_{P_T}^{n_2, n'_2, \nu_2, \nu'_2} r_{T,L}(\nu'_1, \nu'_2). \tag{29}
\end{aligned}$$

This completes the calculation.

Note that at $p_T = 0$, we have

$$I_0^{nn', \nu\nu'} = 2\pi^2 \delta_{nn'} \delta(\nu - \nu') \tag{30}$$

and thus, using (9), we obtain

$$\frac{d\sigma}{dM^2 d^2 p_T} \Big|_{p_T=0} = 8\pi \alpha_{em} \alpha^4 N_c e_f^2 \left(\frac{Qr_0}{2}\right)^2$$

$$\sum_{n_1} \int \frac{d\nu_1}{\pi} e^{\omega(n_1, \nu_1)} Y d_{n_1, \nu_1}(r_0)^{2i\nu_1} \psi_{T,L}^{n_1, \nu_1}(Q, M) \sum_{n_2} \int \frac{d\nu_2}{\pi} e^{\omega(n_2, \nu_2)} Y \bar{d}_{n_2, \nu_2}(r_0)^{-2i\nu_2} \bar{\psi}_{T,L}^{n_2, \nu_2}(Q, M) r_{T,L}(\nu_1, \nu_2), \quad (31)$$

which, for $n_1 = n_2 = 0$, is identical to the result obtained in [8].

3. To obtain more insight into the P_T dependence of the cross-section given by (29), it is useful to evaluate the integrals over ν'_1 and ν'_2 in terms of residues of the relevant poles of the integrands. To this end we observe that the convergence properties of these integrals are determined by the factor

$$\left(\frac{1}{\hat{p}_T} \right)^{2i\nu'} \equiv \left(\frac{4Q}{P_T \sqrt{\beta}} \right)^{2i\nu'}. \quad (32)$$

This factor is easily identified when the explicit expressions (19,22,23) are introduced into the product $\Psi_{T,L}^{n', \nu'}(Q, M) \mathcal{I}_{P_T}^{n', \nu', \nu'}$. Thus for

$$\hat{p}_T \equiv \frac{P_T \sqrt{\beta}}{4Q} < 1, \quad (33)$$

the contour integrals in ν'_1 and ν'_2 must be chosen in the lower half of the complex plane, so only the poles at $\Re(i\nu') \leq 0$ contribute. The integrals are given by the residues of the *moving* poles $i\nu' = \pm i\nu - 2p$ leading to an expansion in terms of powers $(\hat{p}_T)^{4p}$ of the “reduced” transverse momentum(33). Thus one obtains:

$$\frac{d\sigma}{dM^2 d^2 P_T} = 8\pi \alpha_{em} \alpha^4 N_c e_f^2 \left(\frac{Q r_0}{2} \right)^2 \times \sum_{n_1} \int \frac{d\nu_1}{\pi} e^{\omega(n_1, \nu_1)} Y d_{n_1, \nu_1} \bar{E}_{P_T}^{n_1, \nu_1}(r_0) \psi_{T,L}^{n_1, \nu_1}(Q, M) \sum_{n_2} \int \frac{d\nu_2}{\pi} \times e^{\omega(n_2, \nu_2)} Y \bar{d}_{n_2, \nu_2} E_{P_T}^{n_2, \nu_2}(r_0) \bar{\psi}_{T,L}^{n_2, \nu_2}(Q, M) r_{T,L}(\nu_1, \nu_2) \times \left[1 + \mathcal{O}(\hat{p}_T)^4 \right]. \quad (34)$$

On the other hand, for

$$\hat{p}_T \equiv \frac{P_T \sqrt{\beta}}{4Q} > 1, \quad (35)$$

the contour integrals must be closed on the upper half of the complex plane. Consequently, only poles of the Ψ functions at $\Re(i\nu') > 0$ contribute. The

integrals are thus given by *fixed* poles at $i\nu' = \frac{3}{2} + 2p$ and $i\nu' = \frac{5}{2} + 2p$ leading to an expansion independent of ν , resulting in a series expansion in powers of $1/\hat{p}_T$. The two regimes are thus governed by different types of singularities, as was already noticed about vertices of BFKL pomerons [16].

4. We would like to close this note by the following remarks.

(a) The presented calculation extends the results of [8] to non-vanishing momentum transfer and thus -together with [7] where the large mass component was calculated- it completes the derivation of the hard diffractive cross-section in the QCD dipole picture.

(b) The P_T -dependence in Eqs. (29) and (34) may be modified by the proton form factor and possibly other non-perturbative effects. As the same effects would operate also for the large mass component in [7], one may hope that the relative weight of the two components is reasonably well described by our result. A comparison with data should therefore be a significant test of the dipole approach. It would be particularly interesting to compare this new result with the satisfactory description of data obtained in an approximate version of the dipole model [12].

c) Our formulae (29,34) sum over all conformal spin values n allowed by selection rules of the BFKL vertices [16]. At high energies the term with $n = 0$ -corresponding to the so-called “hard pomeron” [1]- dominates. Consequently the phenomenological discussion is usually restricted to $n = 0$. As argued recently [19], however, the analysis of total cross-sections data [20] allows the interpretation of the next term in the expansion, namely $n = 2$, as being at the origin of the so-called “soft pomeron” [21]. Our formula can thus provide an independent test of this hypothesis in hard diffraction dissociation.

To summarize, using the QCD dipole picture of BFKL dynamics we have calculated the quasi elastic component of the diffractive γ^* -dipole cross-section, dominating at small M^2/Q^2 . This work, together with [7], completes the calculation of hard diffraction in the framework of the QCD dipole model and thus allows an extension of the previous phenomenological analyses based on approximate formulae [11, 12]. This should in turn allow to separate perturbative and non perturbative QCD contributions to the diffraction on the proton target. Finally, let us note that the formulae obtained in [7] and here can easily be generalized to $\gamma^* - \gamma^*$ interactions and thus provide a basis for a phenomenological analysis of future collider data.

Acknowledgements

We thank S.Munier and Ch.Royon for fruitful discussions. A.B. thanks the Service de Physique Théorique of Saclay for the kind hospitality. This work was supported in part by the KBN Grant No 2 P03B086 14.

References

- [1] L.N.Lipatov, *Sov. J. Nucl. Phys.* **23** (1976) 642; V.S.Fadin, E.A.Kuraev and L.N.Lipatov, *Phys. Lett.* **B60** (1975) 50; E.A.Kuraev, L.N.Lipatov and V.S.Fadin, *Sov.Phys.JETP* **44** (1976) 45, **45** (1977) 199; I.I.Balitsky and L.N.Lipatov, *Sov.J.Nucl.Phys.* **28** (1978) 822
- [2] J.Bartels, *Zeit. fur. Phys.* **C60** (1993) 471; *Phys. Lett.* **B298** (1993) 204; J.Bartels, M.Wüsthoff *Zeit. fur. Phys.* **C66** (1995) 157; J.Bartels, L.N.Lipatov, M.Wüsthoff, *Nucl. Phys.* **B464** (1996) 298.
- [3] N.N.Nikolaev, B.G.Zakharov, *Zeit. fur. Phys.* **C49** (1991) 607; M.Genovese, N.N.Nikolaev, B.G.Zakharov, *J. Exp. Theor. Phys.* **81** (1995) 625, 633.
For a recent review, see N.N.Nikolaev, *Acta.Phys.Pol.* **B29** (1998) 2425.
- [4] K.Golec-Biernat, J. Kwiecinski, *Phys. Lett.* **B353** (1995) 329. For a recent review see J.Kwiecinski, *Acta.Phys.Pol.* **B27** (1996) 3455.
- [5] S.J. Brodsky et al., *Phys. Rev.* **D50** (1994) 3134; L. Frankfurt, W. Koepf and M. Strikman, *Phys. Rev.* **D54** (1996) 3194. H.Abramowicz, L.Frankfurt and M.Strikman, hep-ph/9503437, *Surveys High Energ. Phys.* **11** (1997) 51.
- [6] A.Bialas and R.Peschanski, *Phys. Lett.* **B378** (1996) 302; **B387** (1996) 405.
- [7] A.Bialas, H.Navelet and R.Peschanski, *Phys. Lett.* **B427** (1998) 147; *Phys. Rev.* **D57** (1998) R6585.
- [8] A.Bialas and W.Czyz, *Acta Phys. Pol.* **B29** (1998) 2095.
- [9] A.H.Mueller, *Nucl. Phys.* **B415** (1994) 373; A.H.Mueller and B.Patel, *Nucl. Phys.* **B425** (1994) 471; A.H.Mueller, *Nucl. Phys.* **B437** (1995) 107.

- [10] For a recent review, see A.H. Mueller, *Acta Phys. Pol.* **B28** (1997) 2557.
- [11] A.Bialas, R.Peschanski and Ch.Royon, *Phys. Rev.* **D57** (1998) 6899.
- [12] S.Munier, R.Peschanski, Ch.Royon, hep-ph/9807488, to appear in *Nucl.Phys.* **B534**.
- [13] H1 Coll., T. Ahmed et al. *Phys. Lett.* **B343** (1995) 681. ZEUS Coll., M.Derrick et al. *Zeit. für. Phys.* **C68** (1995) 569. H1 Coll. C. Adloff et al., Preprint DESY 97-158 (1997).
- [14] ZEUS Coll., M.Derrick et al, *Eur.Phys.J.* **C1** (1998) 81.
- [15] J.D.Bjorken, J.Kogut and Soper, *Phys.Rev.* **D3** (1971) 1382.
- [16] H.Navelet and R.Peschanski, *Nucl.Phys.* **B507** (1997) 353; **B515** (1998) 269.
- [17] L.N. Lipatov *Zh. Eksp. Teor. Fiz.***90** (1986) 1536 (Eng. trans. *Sov. Phys. JETP* **63** (1986) 904).
- [18] I.S.Gradstein and I.M.Ryzhik, *Table of integrals and products*, ed. A.Jeffrey (Acad. Press, N.Y., 1994).
- [19] S.Munier, R.Peschanski, to appear.
- [20] A.Donnachie, P.Landshoff, *Phys. Lett.* **B457** (1998) 408.
- [21] A.Donnachie, P.Landshoff, *Phys. Lett.* **B296** (1992) 227.